

Author Index

Abuin, G., Colli, W., De Souza, W. and Alves, M.J.M.
A surface antigen of *Trypanosoma cruzi* involved in cell invasion (Tc-85) is heterogeneous in expression and molecular constitution, 229

Akella, R., See Vaidya, A.B., 97

Allsopp, B., Carrington, M., Baylis, H., Sohal, S., Dolan, T. and Iams, K.
Improved characterization of *Theileria parva* isolates using the polymerase chain reaction and oligonucleotide probes, 137

Alunda, J.M., see Balaña-Fouce, R., 43

Alves, M.J.M., see Abuin, G., 229

Asikin, N., see Grandea III, A.G., 31

Atiya, A., see Zidovetzki, R., 199

Balaña-Fouce, R., Ordóñez, D. and Alunda, J.M.
Putrescine transport system in *Leishmania infantum* promastigotes, 43

Banerjee, A.B., see Ghoshal, K., 21

Baylis, H., see Allsopp, B., 137

Beach, D.H., Pascal, Jr., R.A. and Holz Jr., GG.
Effects of thiaستearic acids on growth and on dihydrosterculic acid and other phospholipid fatty acyl groups of *Leishmania* promastigotes, 57

Beecroft, R.P., see Tolson, D.L., 109

Boyd, D., De Vos, T., Klassen, G. and Dick, T.
Characterization of the ribosomal DNA from *Trichinella spiralis*, 67

Breier-Saraiva, E.M., see Silva Filho, F.C., 73

Carrington, M., see Allsopp, B., 137

Casper, P., Gentz, R., Matile, H., Pink, J.R. and Sinigaglia, F.
The circumsporozoite protein gene from NF54, a *Plasmodium falciparum* isolate used in malaria vaccine trials (Short Communication), 185

Chambers, A.E., see Harnett, W., 119

Chaudhuri, J. and Donahue, M.J.
Serotonin receptors in the tissues of adult *Ascaris suum*, 191

Cohen, C., see Grandea III, A.G., 31

Colli, W., see Abuin, G., 229

Dalton, J.P. and Heffernan, M.
Thiol proteases released in vitro by *Fasciola hepatica*, 161

Davis, T.B., see Grandea, III, A.G., 31

Davis, W.C., see Reduker, D.W., 239

De Boeck, H., see Zidovetzki, R., 199

De Souza, W., see Abuin, G., 229

De Souza, W., see Silva Filho, F.C., 73

De Souza, W., see Lafaille, J.J., 127

De Vos, T., see Boyd, D., 67

Dick, T., see Boyd, D., 67

Dolan, T., see Allsopp, B., 137

Donahue, M.J., see Chaudhuri, J., 191

Dresden, M.H., see Rege, A.A., 89

Fidge, N.H., see Rogers, M.V., 79

Fuhrman, J.A. and Piessens, W.F.
A stage-specific calcium-binding protein from microfilariae of *Brugia malayi* (Filariidae), 249

Gentz, R., see Caspers, P., 185

Gero, A.M.
Induction of nucleoside transport sites into the host cell membrane of *Babesia bovis* infected erythrocytes, 269

Ghoshal, K., Banerjee, A.B. and Ray, S.
Methylglyoxal-catabolizing enzymes of *Leishmania donovani* promastigotes, 21

Goff, W.L., see Reduker, D.W., 239

Goldenberg, S., see Lafaille, J.J., 127

Grandea III, A.G., Tuyen, L.K., Asikin, N., Davis, T.B., Philipp, M., Cohen, C. and McReynolds, L.A.
A λgt11 cDNA recombinant that encodes *Dirofilaria immitis* paramyosin, 31

Hannon, G.J., see Maroney, P.A., 277

Harnett, W., Chambers, A.E., Renz, A. and Parkhouse, R.M.E.
An oligonucleotide probe specific for *Onchocerca volvulus*, 119

Hedstrom, L. and Wang, C.C.
Purine base transport in wild-type and mycophenolic acid-resistant *Tritrichomonas foetus*, 219

Heffernan, M., see Dalton, J.P., 161

Henkle, K.J., see Rogers, M.V., 79

Herrera, P.R., see Rege, A.A., 89

Higashi, G.I., see Werner, C., 209

Holz, Jr., G.G., see Beach, D.H., 57

Howard, R.J., see Panton, L.J., 149

Iams, K., see Allsopp, B., 137

Jansen, J., see Wesseling, J.G., 167

Jasmer, D.P., see Reduker, D.W., 239

Klassen, G., see Boyd, D., 67

Krieger, M.A., see Lafaille, J.J., 127

Kwakye-Berko, F. and Meshnick, S.R.
Binding of chloroquine to DNA, 51

Lafaille, J.J., Linss, J., Krieger, M.A., Souto-Padrón, T., De Souza, W. and Goldenberg, S.
Structure and expression of two *Trypanosoma cruzi* genes encoding antigenic proteins bearing repetitive epitopes, 127

Leech, J.H., see Rosenthal, P.J., 177

Linss, J., see Lafaille, J.J., 127

Lopez, M., see Rege, A.A., 89

Maloy, W.L., see Panton, L.J., 149

Maroney, P.A., Hannon, G.J. and Nilsen, T.W.
Accurate and efficient RNA polymerase III transcription in

a cell-free extract prepared from *Ascaris suum* embryos, 277

Matile, H., see Caspers, P., 185

McGuire, T.C., see Reduker, D.W., 239

McKerrow, J.H., see Rosenthal, P.J., 177

McPhie, P., see Panton, L.J., 149

McReynolds, L.A., see Grandea III, A.G., 31

Meshnick, S.R., see Kwakye-Berko, F., 51

Mitchell, G.F., see Rogers, M.V., 79

Myler, P.J., see Scholler, J.K., 11

Nilsen, T.W., see Maroney, P.A., 277

Ordóñez, D., see Balaña-Fouce, R., 43

Panton, L.J., McPhie, P., Maloy, W.L., Wellem, T.E., Taylor, D.W. and Howard, R.J.
Purification and partial characterization of an unusual protein of *Plasmodium falciparum*: histidine-rich protein II, 149

Parkhouse, R.M.E., see Harnett, W., 119

Pascal Jr., R.A., see Beach, D.H., 57

Pearson, T.W., see Tolson, D.L., 109

Perryman, L.E., see Reduker, D.W., 239

Philipp, M., see Grandea III, A.G., 31

Piessens, W.F., see Fuhrman, J.A., 249

Pink, J.R., see Caspers, P., 185

Rajan, T.V., see Werner, C., 209

Rasnick, D., see Rosenthal, P.J., 177

Ray, S., see Ghoshal, K., 21

Reduker, D.W., Jasmer, D.P., Goff, W.L., Perryman, L.E., Davis, W.C. and McGuire, T.C.
A recombinant surface protein of *Babesia bovis* elicits bovine antibodies that react with live merozoites, 239

Rege, A.A., Herrera, P.R., Lopez, M. and Dresden, M.H.
Isolation and characterization of a cystein proteinase from *Fasciola hepatica* adult worms, 89

Remold, H., see Toye, P., 1

Renz, A., see Harnett, W., 119

Rogers, M.V., Henkle, K.J., Fidge, N.H. and Mitchell, G.F.
Identification of multispecific lipoprotein receptor in adult *Schistosoma japonicum* by ligand blotting analyses, 79

Rosenthal, P.J., McKerrow, J.H., Rasnick, D. and Leech, J.H.
Plasmodium falciparum: inhibitors of lysosomal cysteine proteinases inhibit a trophozoite proteinase and block parasite development, 177

Schoenmakers, J.G.G., see Wesseling, J.G., 167

Scholler, J.J., Myler, P.J. and Stuart, K.D.
A novel telomeric gene conversion in *Trypanosoma brucei*, 11

Sherman, I.W., see Zidovetzki, R., 199

Silva Filho, F.C., Breier-Saraiva, E.M., Tosta, M.X. and De Souza, W.
Trichomonas vaginalis and *Tritrichomonas foetus* secrete neuraminidase into the culture medium, 73

Sinigaglia, F., see Caspers, P., 185

Smits, M.A., see Wesseling, J.G., 167

Snijders, P.J.F., see Wesseling, J.G., 167

Sohal, S., see Allsopp, B., 137

Souto-Padrón, T., see Lafaille, J.J., 127

Stuart, K.D., see Scholler, J.K., 11

Suplick, K., see Vaidya, A.B., 97

Swamy, K.H.S.,
Filarial parasites exhibit unusually high levels of choline acetyltransferase activity, 259

Taylor, D.W., see Panton, L.J., 149

Tolson, D.L., Turco, S.J., Beecroft, R.P. and Pearson, T.W.
The immunochemical structure and surface arrangement of *Leishmania donovani* lipophosphoglycan determined using monoclonal antibodies, 109

Tosta, M.X., see Silva Filho, F.C., 73

Toye, P. and Remold, H.
The influence of temperature and serum deprivation on the synthesis of heat-shock proteins and alpha and beta tubulin in promastigotes of *Leishmania major*, 1

Turco, S.J., see Tolson, D.L., 109

Tuyen, L.K., see Grandea III, A.G., 31

Vaidya, A.B., Akella, R. and Suplick, K.
Sequences similar to genes for two mitochondrial proteins and portions of ribosomal RNA in tandemly arrayed 6-kilobase-pair DNA of a malarial parasite, 97

Van Someren, P., see Wesseling, J.G., 167

Wang, C.C., see Hedstrom, L., 219

Wellem, T.E., see Panton, L.J., 149

Werner, C., Higashi, G.I., Yates, J.A. and Rajan, T.V.
Differential recognition of two cloned *Brugia malayi* antigens by antibody class, 209

Wesseling, J.G., Snijders, P.J.F., Van Someren, P., Jansen, J., Smits, M.A. and Schoenmakers, J.G.G.
Stage-specific expression and genomic organization of the actin genes of the malaria parasite *Plasmodium falciparum*, 167

Yates, J.A., see Werner, C., 209

Zidovetzki, R., Sherman, I.W., Atiya, A. and De Boeck, H.
A nuclear magnetic resonance study of the interactions of the antimalarials chloroquine, quinacrine, quinine and mefloquine with dipalmitoylphosphatidylcholine bilayers, 199

Subject Index

Acetylcholine, 259
 Actin gene, 167
 Adenosine, 269
 Affinity-purified antibody, 31
 Altered nucleoside permeation, 269
 Antigen, 127
 Antigenic variation, 11
 Antimalarial, 51
 Antimalarial drug, 149
 Apoprotein, 79
Ascaris suum, 191, 277

Babesia bovis, 239, 269
 Berenil, 43
 Bovine erythrocyte, 269
Brugia malayi, 209, 249, 259, 277
Brugia pahangi, 259

 Calcium-binding protein, 249
 Cell invasion, 229
 Chagas' disease, 127
 Chemotherapy, 177
 Chloroquine, 51
 Choline acetyltransferase, 259
 Circumsporozoite protein gene, 185
 Collagen, 209
 Cysteine proteinase, 89
 Cytochrome *b*, 97
 Cytochrome oxidase, 97
 Cytotoxicity, 73

 α -Difluoromethylornithine, 43
 Dihydrosterculic acid, 47
Dirofilaria immitis, 31
 DNA, 51
 rDNA, 67
 DNA probe, 119

 Ethacrynic acid, 259
 Evolution, 97

Fasciola hepatica, 89, 161
 Fatty acyl group, 57
 Filariiae, 259
 Fluoromethylketone, 177
 Fusion protein, 239

 Gene cloning, 127
 Gene conversion, 11
 Glycoprotein, 229

 Haemoglobin, 177
 Heat-shock protein, 1
 Histidine-rich protein, 149
 Hypoxanthine, 219

 IgE, 209
 IgG, 209
 In vitro released product, 161
 In vitro transcription, 277
 Intercalator, 51
 Intergenic spacer, 67
 Internal transcribed spacer, 67
 Intestine, 191
 Intron, 167

 D-Lactate, 21
Leishmania, 57
Leishmania donovani, 21, 109
Leishmania donovani infantum, 43
Leishmania infantum, 43
Leishmania major, 1, 109
 Ligand blotting, 79
 Lipid membrane, 199
 Lipophosphoglycan, 109
 Lipoprotein receptor, 79
 [125 I]Lipoprotein, 79
Litomosoides carinii, 259
 Lymphatic filariasis, 209, 259
 λ gt11 cDNA library, 31
 λ gt11 fusion protein expression, 209
 λ gt11 genomic library, 239

 Malaria, 51, 177, 185
 Methylglyoxal, 21
 Microfilariae, 249
 Mitochondrial DNA, 97
 Monoclonal antibody, 109, 239, 249
 Muscle, 191
 Mycophenolic acid resistance, 219
 Myosin, 31, 209

 Neuraminidase, 73
 Nitrobenzylthioinosine, 269
 Non-coding region, 167
 Nuclear magnetic resonance, 199
 Nucleoside transport, 269

 Oligonucleotide probe, 137
Onchocerca volvulus, 119

 Paramyosin, 31
 Parasitic nematode, 191
 Parasitic protozoon, 73
 Phosphatidylcholine, 57
 Phosphatidylethanolamine, 57
Plasmodium, 51, 97
Plasmodium falciparum, 149, 167, 177, 185, 269
 Polyamine, 43
 Polymerase chain reaction, 137
 Promastigote, 57, 109

Protein structure, 149
Protease inhibitor, 161
Proteinase, 177
Protozoa, 21
Purine transport, 219
Putrescine transport, 43

Recombinant protein, 239
Repeat unit, 67
Repetitive epitope, 127
Restriction mapping, 67
Ribosomal RNA, 97
RNA polymerase III, 277
5S rRNA, 277

Schistosoma japonicum, 79
Schistosoma mansoni, 79
Serotonin receptor, 191
Speciation, 119
Stage-specific transcription, 167
Strain characterization, 137
Surface epitope, 109

Surface heterogeneity, 229
Surface-exposed epitope, 239

Telomere, 11
Theileria parva, 137
Thiastearic acid, 57
Thiol protease, 161
Transmission-blocking immunity, 249
Trematode, 161
Trematode, parasitic, 89
Trichinella spiralis, 67
Trichomonas vaginalis, 73
Tritrichomonas foetus, 73, 219
Trypanosoma brucei, 11
Trypanosoma cruzi, 127, 229
Tubulin, 1

Worm motility, 259
Wuchereria bancrofti, 259

Zinc-chelate chromatography, 149



